RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms

Toxic to fish and aquatic organisms. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's Certification



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF KANSAS

EPA Reg. No. 100-1295 EPA SLN No. KS-130003

For Control of Alfalfa Weevil Infesting Alfalfa

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

These directions for use must be in the possession of the user at the time of application.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restriction, and precautions on the EPA-registered product label for Warrior II with Zeon Technology (EPA Reg. No. 100-1295) are to be followed.

		Rate	
Crop	Target Pests	lb. aid./A	fl. oz./A
ALFALFA AND ALFALFA GROWN FOR SEED	Alfalfa Weevil Larvae and Adults	0.02 - 0.03	1.28 – 1.92

Foliar Applications

Maximum Application Rate: If no more than 0.1 lb a.i. (6.25 fl. oz. or 0.4 pts. of product) has been applied during the annual alfalfa growing season, Warrior II with Zeon Technology may be applied at 1.28 - 1.92 fl. oz. per acre after the last annual cutting and before alfalfa and insects become dormant to help reduce overwintering populations of alfalfa insect pests and subsequent insect populations the following spring. Treatments in this management program are part of the total annual maximum active ingredient allowed per acre per season.

- Apply Warrior II with Zeon Technology as a broadcast foliar spray with either ground or aerial application equipment.
- Use a minimum of 10 GPA by ground or 2 GPA by aerial application. When foliage is dense and/or pest populations are high 5–10 gals. per acre by air or 20 gals. per acre by ground and higher use rates are recommended. Use higher rates for increased residual control.
- Warrior II with Zeon Technology is labeled for chemigation application. Refer to Chemigation use directions on the full Warrior II with Zeon Technology label for details on this application.
- Make applications when insect populations reach economic thresholds or when significant feeding is occurring.
- This product is highly toxic to bees exposed to direct treatment. Avoid application when bees are actively foraging by applying during the early morning or during the evening hours. Be aware of bee hazard resulting from a cool evening and/or morning dew. It may be advisable to remove bee shelters during and for 2–3 days following application. Avoid direct application to bee shelters.
- Observe all spray drift precautions on the EPA registered label.
- Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results.

RESTRICTIONS:

- Do not apply within 1 day of harvest for forage or within 7 days of harvest for hay or seed.
- Do not make applications of Warrior II with Zeon Technology or follow with other products containing lambda-cyhalothrin less than 7 days apart.
- Do not exceed 7.68 fl. oz. /A (0.12 lbs. a.i.) or make more than 4 applications of Warrior II with Zeon Technology at 1.92 fl. oz./A (0.03 lbs. a.i.) or other products containing lambda-cyhalothrin per season.
- Re-entry Interval: 24 hours. Personal protective equipment is required for early entry.

WARRIOR II with Zeon Technology® trademark of a Syngenta Group Company

Registrant: Syngenta Crop Protection, LLC PO Box 19300 Greensboro, NC 27419

KS1295028AA1113