

## **RESTRICTED USE PESTICIDE**

DUE TO HIGH ACUTE INHALATION TOXICITY AND EYE AND SKIN CORROSIVENESS TO HUMANS; HIGH ACUTE TOXICITY TO NONTARGET BIRDS AND AQUATIC INVERTEBRATES, AND THE NEED FOR HIGHLY SPECIALIZED APPLICATOR TRAINING.

For retail sale to, and use only by, USDA APHIS Certified Applicators trained in bird control or by persons under their direct supervision.

### SUPPLEMENTAL LABELING

## **COMPOUND DRC-1339 CONCENTRATE STAGING AREAS – KANSAS EPA Registration Number 56228-30**

SLN KS-120003

### **FOR DISTRIBUTION AND USE ONLY WITHIN KANSAS**

This product may be used to control common crows (*Corvus brachyrhynchos*) and fish crows (*Corvus ossifragus*) in non-crop areas surrounding peanut fields and pecan orchards or in bait containers within peanut fields and pecan orchards during September, October, November, December, January and February only.

**THIS LABEL AND THE FEDERAL LABEL MUST BE IN POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION. READ THE ENTIRE LABEL AND THE ENTIRE FEDERAL LABEL. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS FOUND ON EITHER LABEL.**

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read the entire label and the label for EPA Registration Number 56228-30. This product must be used strictly in accordance with both labels' precautionary statements and applicable use directions, as well as with all applicable State and Federal laws and regulations.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable State wildlife agency and obtain all necessary kill or collecting permits.

**GENERAL INFORMATION:** This product contains a slow acting avicide which kills target birds in 1 to 3 days. As many types of nontarget bird species are potentially vulnerable to DRC-1339, it is necessary to use care and to follow the requirements of this label to minimize impacts to non-target species.

**USE RESTRICTIONS:** Baits made from Compound DRC-1339 Concentrate-KS may only be used to control common crows (*Corvus brachyrhynchos*) and fish crows (*C. ossifragus*) in noncrop areas surrounding peanut fields and pecan orchards or in bait containers within peanut fields and pecan orchards and only during September, October, November, December, January and February. Do not use baits made from this product from March 1<sup>st</sup> through August 31<sup>st</sup> of any year.

- **DO NOT** apply baits made from this product by air or using and procedures not prescribed on this label.
- This product may only be used to make baits with the bait materials listed on the Federal label and rolled corn, rolled barley, brown rice, poultry pellets, whole corn, cull or waste potatoes, dry dog food, shelled pecans, or shelled peanuts following the instructions indicated on this label under **BAIT PREPARATION**.
- **DO NOT** use baits as food or feed.
- **DO NOT** apply this product in a manner that will contaminate food, feed, or water.
- **DO NOT** harvest any crop for use as food or feed that has been contaminated with this product.
- **DO NOT** graze livestock or plant any rotational crop in treated noncrop areas or in crop areas that may have been contaminated by treated bait spilled from bait application containers for one year (365 days) following the last application of bait made from this product.
- **DO NOT exceed a maximum single application rate of 0.09 lbs of active ingredient per treated acre or a maximum cumulative application rate of 0.5 lbs of active ingredient per treated acre in any use year.** The maximum single treatment rate of 0.09 lbs of active ingredient per acre would be provided by one treatment with a 2% DRC-1339 bait diluted 1:10 with untreated bait and applied at a rate of 52 lbs of diluted bait per acre.
- **DO NOT** apply treated baits within 50 feet of bodies of water, unless baited sites are under constant observations while baits are exposed.
- Before baits made from this product are applied, sites that are to be treated must be observed for evidence of nontarget activity and must be prebaited (see specific instructions for these activities). **DO NOT** apply treated baits in locations where prebait has not been accepted well by nontarget species or where nontarget wildlife have been observed to feed on prebait.
- **DO NOT** apply bait(s) in areas where there is danger that Threatened or Endangered species will consume baits unless special precautions are taken to limit such exposures. Such precautions shall include observation of baited sites and hazing tactics to frighten away Threatened or Endangered species that otherwise might feed upon baits. A permit may be required for harassment of Threatened or Endangered Species. See **THREATENED AND ENDANGERED SPECIES** section on the product label for EPA Reg. No. 56228-30 for information on consideration of threatened and endangered species.
- **DO NOT** store, apply, or even temporarily place treated bait in locations accessible to children, pets, or domestic animals or non-target animals.
- **DO NOT** apply baits made from this product in a way that will contact workers or other persons.

**ENTRY RESTRICTIONS:** Keep persons other than handlers, as well as pets and livestock, away from the bait at all times. Only protected handlers may be in the area during bait application.

**PRETREATMENT OBSERVATIONS:** Prior to application of treated bait, observe target populations thoroughly to determine their relative numbers, feeding habits, preferred staging areas or feeding locations, daily activity patterns, and the relative abundance and types of nontarget species at potential baiting sites associated with pecan orchards. Once potential baiting sites are identified, they must be observed to determine the potential hazards to nontarget species and prebaited to promote bait acceptance by target species. **DO NOT** apply treated bait where there is a danger that it will be consumed by threatened or endangered species.

**PREBAITING:** After daily activity patterns of birds have been noted and appropriate baiting location has been selected, apply untreated bait material of the type that will be used for toxic baiting at the same site. Apply prebait using the application method and rate that is to be used initially for toxic bait (see **APPLICATION DIRECTIONS**), making sure that prebait used in peanut fields or pecan orchards is placed in bait application containers.

**BAIT PREPARATION:** Bait batch size may be scaled up or down to prevent waste and accommodate mixing equipment.

**CRACKED CORN, ROLLED BARLEY AND BROWN RICE BAITS**

Before mixing baits, screen brown rice on a Y Grade sieve to remove small particles and dust and to retain rice particles weighing between 20 and 40 mg each (roughly 11,000 to 23,000 particles per lb).

- Place 10 lbs (4.5 kg) of the grain selected into a mixer or a plastic container or a leak-proof plastic bag and set aside.
- Warm a mixture of 10 ml (0.3 oz) of edible oil and 30 ml (1 oz) of Alcolec-S (lecithin) to 110 °F (43 °C) for about 5 minutes. Dissolve 92 grams (3.2 oz) of Compound DRC-1339 Concentrate-KS in the mixture of edible oil and lecithin.
- Pour resulting solution or suspension over the 10 lbs of selected grain material and mix or tumble slowly until the bait is evenly mixed.
- Air dry bait at ambient temperature. When bait is dry, mix treated bait with untreated bait material of the same type selected for the toxic bait at a ratio from one part of treated bait to 10 to 25 parts of untreated bait.

**POULTRY PELLETS AND WHOLE CORN BAITS**

- Place 10 lbs (4.5 kg) of the poultry pellets or whole corn into a mixer or a plastic container or a leak-proof plastic bag and set aside.
- Warm 400 ml (14 oz) of potable water to 110 °F (43 °C) for about 5 minutes. Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate-KS in the warmed water.

- Pour the resulting solution or suspension over the 10 lbs of poultry pellets or whole corn and mix or tumble slowly until the bait is evenly mixed.
- Air dry bait at ambient temperature. When bait is dry, mix treated bait with untreated bait material of the same type selected for the toxic bait at a ratio of one part treated bait to 10 to 25 parts of untreated bait.

#### **CULL OR WASTE POTATO BAITS**

- Place 50 lbs. (23 kg) of cull or waste potatoes in a mixer or a plastic container or a leak-proof plastic bag and set aside.
- Warm 600 ml (1.3 pt) of potable water to 110 °F (43 °C) for about 5 minutes. Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate-KS in the warmed water.
- Pour the resulting solution or suspension over the 50 lbs. of cull or waste potatoes and mix or tumble slowly until the bait is evenly mixed.
- Air dry bait at ambient temperature. When bait is dry, mix treated bait with untreated bait material of the same type selected for the toxic bait at a ratio of no less than 1 part treated bait to 2 parts untreated bait.

#### **DRY DOG FOOD BAITS**

- Place 10 lbs (4.5 kg) dry dog food in a plastic container, plastic drop cloth, or mixer and set aside.
- Warm 600 ml (1.3 pt) of potable water to 110 °F (43 °C) for about 5 minutes. Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate-KS in the warmed water. Binding agents, such as lecithin oils or polyvinyl acetates, may be used to enhance retention and to reduce losses due to moisture when mixing DRC-1339 Concentrate-KS with water. Do not use more than 3 oz (85 g) binding agent per 10 lb of bait.
- If desired, 46 grams (1.6 oz) of Compound DRC-1339 Concentrate-KS can be dissolved in 280 ml (10 oz) of edible oil instead of warm potable water.
- Pour the resulting solution or suspension over the 10 lb of dry dog food and mix or tumble until bait is evenly distributed.
- Air dry bait at ambient temperature. When bait is dry, dilute treated bait with untreated bait material of the same type selected for the toxic bait at a rate of at least 1 part treated bait to 5 parts untreated bait.

#### **SHELLED PEANUT OR SHELLED PECAN BAITs**

- Place 10 lbs (4.5 kg) of shelled peanuts or shelled pecans in a mixer or a plastic container or leak-proof plastic bag and set aside.
- Warm 400 ml (14 oz) of potable water to 110°F (43°C) for about 5 minutes. Dissolve 46 grams (1.6 oz) of Compound DRC-1339 Concentrate-KS in the warmed water.
- Pour the resulting solution or suspension over the 10 lbs of shelled peanuts or shelled pecans and mix or tumble slowly until the bait is evenly distributed.
- Air-dry bait at ambient temperature. When bait is dry, mix treated bait with untreated bait material of the same type selected for the toxic bait at a ratio from one part of treated bait to 10 to 25 parts of untreated bait.

Treated baits prepared as described above and stored under cool and dry conditions may be used for up to 30 days after preparation. Use diluted baits within 7 days after preparation. Immediately use treated or diluted baits that have been exposed to sunlight or heat in excess of 110 °F (43 °C) following preparation, or dispose of them as pesticide waste.

**APPLICATION DIRECTIONS:** Wear rubber gloves while handling treated baits and bait mixtures and dispense or broadcast baits using a scoop or other appropriate equipment.

**MANUAL BROADCAST APPLICATIONS (NON-CROP AREAS ONLY):** When target birds are observed feeding well on the prebait and about 75% of the prebait has been consumed apply treated baits (diluted 1:10 to 1:25 with untreated bait as described above under BAIT PREPARATION). Scatter treated bait manually or by use of a mechanical spreader within the non-crop area at a rate of 10 to 25 lbs diluted bait per acre (11 to 28 kg/hectare).

The baiting rate maybe increased to 52 lbs per acre (58 kg/hectare) of diluted bait, **if warranted** by feeding pressure from target species. Re-treat when 75% of the previously applied treated baits have been consumed, following 0.4 inches (10 mm) or more of rain, or at 4-day intervals for as long as target species continue to visit the treated area. Do not exceed a treatment rate per application of 0.09 lbs of the active ingredient per acre or a cumulative treatment rate of 0.5 lbs of active ingredient per acre for the use year.

Cease baiting if target species are observed to feed on, or are adversely affected by, the treated bait.

**BAIT DISPENSER APPLICATIONS:** Baits applied within peanut fields or pecan orchards must be placed in bait dispensers which minimize the possibility of spillage. Position bait dispensers in a manner that encourages feeding by target crows and prevents contamination of food and feed crops or water due to spillage of bait.

Replenish treated bait when 75% or more of the previously offered bait has been consumed. The total amount of active ingredient used to bait pecan orchards or peanut fields may not exceed 0.5 lbs of active ingredient per acre.

- **DO NOT** bait in areas that are occupied or grazed by livestock. Cease baiting if non-target species are observed to feed on, or are adversely affected by, treated bait.

**POSTTREATMENT CLEANUP:** Remove all unconsumed bait and any spilled bait at the conclusion of the treatment period. Dispose of collected, unused, and outdated bait by burial in an approved sanitary landfill in accordance with Federal and State laws or by incineration at an approved site if permitted by State and local authorities.

Search treated areas, animal pens, and immediate surrounding areas for bird carcasses. Collect carcasses of dead or dying birds and bury or burn them according to applicable laws. Repeat carcass searches at 1-2 day intervals as long as bait is exposed and likely to remain toxic, and after baiting has stopped.

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store only in original container, in a dry place inaccessible to children, pets, and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bags by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. If bags are not to be recycled, dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.

**EPA SLN NUMBER KS-120003**

**24(c) Registrant:**

U.S. Department of Agriculture  
Animal and Plant Health Inspection Service  
4700 River Road, Unit 149  
Riverdale, MD 20737-1237

**Local Contact:**

State Director  
USDA APHIS  
Wildlife Services  
4070 Ft. Riley Blvd  
Manhattan, KS 66502  
(785) 537-6855

Batch Code No.: \_\_\_\_\_

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND  
DOMESTIC ANIMALS  
DANGER**

**Acute Hazards:** Fatal if inhaled. Corrosive. Causes irreversible eye damage and skin burns. May be fatal if swallowed. Harmful if absorbed through skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some people.

**Hazard Avoidance:** Do not get in eyes, on skin, or on clothing. Do not breathe dust. Wear protective clothing, protective eyewear, and respiratory protection as listed under "Personal Protective Equipment." Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. Remove contaminated clothing and wash before reuse.

**PERSONAL PROTECTIVE EQUIPMENT:**

- Handlers who mix packages containing 1 lb or more of this product must wear:**
- Coveralls over long-sleeved shirt and long pants
  - Chemical-resistant gloves (such as waterproof or rubber gloves)
  - Chemical-resistant footwear plus socks
  - Protective eyewear (goggles or face shield)
  - Respirator with a dust/mist filtering respirator (MSHA NIOSH approval number prefix TC-21C or NIOSH approved respirator with any N, R, P, or HE filter)

**Handlers who mix packages containing less than 1 lb of this product must wear:**

- Chemical-resistant gloves (such as waterproof or rubber gloves)
- Protective eyewear (goggles or face shield)

**Applicators who handle bait must wear:**

- Chemical-resistant gloves (such as waterproof or rubber gloves)

**RESTRICTED USE PESTICIDE**

Due to High Acute Inhalation Toxicity, Eye and Skin Corrosiveness to Humans; High Acute Toxicity to Nontarget Birds and Aquatic Invertebrates; and the Need for Highly Specialized Applicator Training. For retail sale to, and use only by USDA APHIS Certified Applicators trained in bird control or by persons under their direct supervision.

**COMPOUND DRC-1339  
CONCENTRATE - STAGING AREAS**

For bird control in non-crop staging areas associated with roosts

Active ingredient  
Stillicide, 3-chloro-p-toluene hydrochloride: 97.0%  
Inert ingredients: 3.0%  
TOTAL: 100.0%

**KEEP OUT OF REACH OF CHILDREN  
DANGER-PELIGRO  
POISON**



**FIRST AID**

Have label with you when obtaining treatment advice.

If swallowed	- Call a poison control center or doctor immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by the poison control center or doctor.
If on skin or clothing	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor immediately for treatment advice.
If inhaled	- Move person to fresh air. - If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. - Call a poison control center or doctor immediately for treatment advice.
If in eyes	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call poison control center or doctor immediately for treatment advice.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage. See additional PRECAUTIONARY STATEMENTS on Left Panel.

**User Safety Requirements:**

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are provided for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:**

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is very highly toxic to birds and aquatic invertebrates. Do not use in any manner that may endanger desirable and protected bird species. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply when runoff is likely to occur. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by the cleaning of equipment or disposal of equipment washwaters or rinsate.

**ENDANGERED SPECIES CONSIDERATIONS**

Before undertaking any control operations with the product, consult with local, State, and Federal Wildlife authorities to ensure the use of this product presents no hazard to any Threatened or Endangered species. DO NOT apply in areas where the product may be consumed by Threatened or Endangered species.

**UNITED STATES DEPARTMENT OF AGRICULTURE  
ANIMAL AND PLANT HEALTH INSPECTION SERVICE**

4700 River Road, Unit 149  
Riverdale, MD 20737  
EPA Reg. No. 56228-30  
California Reg. No. 56228-30-AA  
EPA Est. No. 56228-ID-1  
Net Contents: \_\_\_\_\_  
Batch Code No.: \_\_\_\_\_



## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### READ THIS LABEL:

Read the entire label. This product must be used strictly in accordance with precautionary statements, use directions, as well as with all applicable State and Federal laws and regulations.

Before using this product, contact the U.S. Fish and Wildlife Service and the applicable wildlife agency and obtain all necessary kill or collecting permits.

### GENERAL INFORMATION:

This product contains a slow-acting avicide which kills target bird species (see lists below) in 1 to 3 days. As many types of nontarget bird species are potentially vulnerable to DRC-1339, it is necessary to use care and to follow the requirements of this label to minimize impacts to nontarget species.

### USE RESTRICTIONS:

Baits made from this product may only be used in noncrop "staging areas" associated with nighttime roosting sites and may only be used to control the types of blackbirds, cowbirds, grackles, and starlings that are specifically listed on this label. Staging areas where baits may be applied are limited to stubble fields, harvested dormant hay fields, open grassy or bare-ground noncrop areas, roads, roadsides, rooftops, industrial and commercial structures, and secured parking areas. Baits must be prepared and applied as specified on this label. Baits made from this product must be prepared with one of the following grain components: brown rice, rolled barley, cracked corn, or poultry pellets. **DO NOT** apply baits made from this product by air. **DO NOT** apply baits made from this product to orchards.

Users of this product must follow all limitations on this label regarding the application of treated baits in staging areas. **DO NOT** use treated baits as food or feed. **DO NOT** graze animals on treated areas for 365 days following the last bait application. **DO NOT** apply treated baits in any way that could contaminate food or feed crops, or that would allow bait to be consumed by livestock. **DO NOT** apply treated baits within 50 feet of bodies of water.

## DIRECTIONS FOR USE (continued)

Before baits made from this product are applied, sites that are to be treated must be observed for evidence of nontarget activity and prebaited (see specific instructions for these activities). **DO NOT** apply baits in areas where there is danger of consumption by Threatened or Endangered species.

**DO NOT** exceed a maximum application rate of 0.1 lb of active ingredient per treated acre per treatment or a maximum yearly application rate of 0.5 lb of active ingredient per acre. An application rate of 0.1 lb of active ingredient per acre is consistent with use of a barley or a cracked corn bait that is 1.9% active ingredient, diluted 1:10 with untreated baits, and applied at a rate of 58 lbs of diluted bait per treated acre. That rate is also consistent with use of a brown rice bait that is 1.9% active ingredient, diluted 1:25 with untreated bait, and applied at a rate of 137 lbs of diluted bait per treated acre (See FORMULATION DIRECTIONS and APPLICATION DIRECTIONS).

Do not apply bait in a way that will contact workers or other persons.

**Rotational Crop (Plantback) Restrictions:** After last application of bait, the plantback intervals are: rice, wheat, corn, and barley (15 days); sunflower and soybeans (30 days); and other crops (365 days).

### ENTRY RESTRICTIONS:

Keep persons other than authorized handlers, as well as pets and livestock, away from the bait at all times. Only protected handlers may be in the area during bait application.

#### Target Species:

Boat-tailed grackle (*Quiscalus major*)  
Brown-headed cowbird (*Molothrus ater*)  
Common grackle (*Quiscalus quiscula*)  
Great-tailed grackle (*Quiscalus mexicanus*)  
Red-winged blackbird (*Agelaius phoeniceus*)  
European Starling (*Sturnus vulgaris*)

## DIRECTIONS FOR USE (continued)

When mixed in flocks with one or more of the species listed above, the following species also shall be considered to be target species: common crow (*Corvus brachyrhynchos*), black-billed magpie (*Pica pica*), Brewer's blackbird (*Euphagus cyanocephalus*), tricolored blackbird (*Agelaius tricolor*), and yellow-headed blackbird (*Xanthocephalus xanthocephalus*).

### PRETREATMENT OBSERVATIONS:

Prior to application of treated bait to any problem area, carefully observe habits of target birds to (1) determine their relative numbers, (2) locate the staging areas which can be baited in accordance with this label, (3) determine the optimum time of day for applying bait, and (4) evaluate the potential for hazards of the application to desirable and protected species.

### SELECTING BAIT MATERIALS:

Use only cracked corn, rolled barley, brown rice, or poultry pellets as the grain component of the bait. Poultry pellets and cracked corn generally are accepted by starlings. Cracked corn, brown rice, and rolled barley generally are accepted by blackbirds.

If you are uncertain as to which bait material to select, expose the target population to small amounts of two or more of the following: cracked corn, rolled barley, brown rice, or poultry pellets to determine a suitable bait material.

## DIRECTIONS FOR USE (continued)

### PREBAITING:

Prebaiting with untreated bait material of the type to be used to toxic baiting is required. Apply prebait at the locations that are to be used for toxic baiting using the same application method and application rate that is to be used for toxic baiting (see APPLICATION DIRECTIONS).

Expose prebait for 3-7 days or until prebait is generally well accepted. Change prebaiting locations within staging area if necessary to achieve good acceptance by target species or if nontarget species have been observed eating the prebait. **DO NOT** apply treated bait at sites where prebait has not been well accepted by target species or where nontarget species have been observed eating prebait.

### BAIT PREPARATION:

**Screening:** Prepare all grain and pelleted baits for treatment by screening through a commercial sieve of grade "Y" (hole size 6/64 inch round) to remove small particles and dust.

**Mixing Bait:** Prepare baits in the amounts shown below:

**Cracked Corn or Rolled Barley (1.9% bait) -**  
92 g (3.2 oz.) Compound DRC-1339 Concentrate  
400 ml (14 oz.) warm potable water

**4.5 Kg (10 lbs.) rolled barley**  
Add up to 30 ml (1 oz) of a binding agent (lecithin oil, latex, or polyvinyl acetate) if needed

**Brown Rice (1.9% bait) -**  
92 g (3.2 oz.) Compound DRC-1339 Concentrate  
10 ml (0.3 oz.) edible oil  
30 ml (1 oz) Alcolac-S  
4.5 kg (10 lbs) brown rice

**Poultry Pellets (0.9% - 1% bait) -**  
46 g (1.6 oz.) Compound DRC-1339 Concentrate  
400 ml (14 oz.) warm potable water  
4.5 kg (10 lbs) poultry pellets

Prepare baits as follows:

1. Dissolve or suspend Compound DRC-1339 Concentrate - Staging Areas in warm potable water or edible oil warmed to 110°F (43°C);
2. Add binding agents as indicated above for the bait formulation selected;
3. Pour the liquid mixture over the bait material selected;
4. Mix or tumble slowly until liquid mixture is distributed evenly over the bait; and
5. Air dry bait at ambient temperature.

## DIRECTIONS FOR USE (Continued)

### BAIT STORAGE:

If treated baits are stored under cool and dry conditions, they may be used for up to 30 days after preparation. Use diluted baits (see below) within 7 days after preparation. Immediately use treated or diluted baits that have been exposed to sunlight or heat in excess of 110°F (43°C) following preparation, or dispose of them as pesticide waste.

### DILUTING BAIT:

Dilute treated baits with untreated, similar grain materials to reduce the possibility that any nontarget bird will ingest more than one toxic particle and to allow more individuals of the target species to consume toxic bait particles. For the baits that may be prepared using this product, the dilution ratios shown below are required.

Bait Material	Ratio of Treated to Untreated Particles
rolled barley, cracked corn, or poultry pellets	1 treated to 10 or more untreated particles
brown rice	1 treated to 25 or more untreated particles

### APPLICATION DIRECTIONS:

Wear rubber gloves when handling bait. Use a scoop or other appropriate utensil when manually applying or loading bait application equipment. Apply bait according to one of the methods described below.

#### 1. Feeding Stations

Place up to 10 lbs in one or more retrievable feeding stations. Place feeding stations in locations within staging areas which appear most likely to selectively attract target species in large numbers. Use the area of the baited portion of the staging area to calculate the per-acre rate. If only one feeding station is used or if stations are more than 140 yards apart, count one acre of treated area for each station used.

Replace diluted bait if 75% or more of the amount originally placed has been consumed, if 0.4 inches (10 mm) of precipitation has fallen, or after 7 days of exposure to birds. Do not replenish bait if target species are no longer present or are no longer feeding on bait. Do not exceed the application rate limits specified below.

## DIRECTIONS FOR USE (Continued)

#### 2. Mechanical Broadcasting

Broadcast bait using ground-based equipment calibrated so as not to exceed the application rate limits specified below. Apply bait to (alternate) swaths 20-50 feet wide that are spaced at least one swath width (20-50 feet) apart. Do not apply bait by use of aircraft.

Re-treat the same swaths or treat previously untreated swaths if 75% or more of the amount originally placed has been consumed, if 0.4 inches (10 mm) of precipitation has fallen, or after 7 days of exposure to birds. Do not replenish bait if target species are no longer present or are no longer feeding on bait. Do not exceed the application rate limits specified below.

#### 3. Manual Broadcasting

Wearing rubber gloves, manually broadcast at a rate suitable for the level of target bird pressure but not exceeding the application rate limits specified below. Apply bait to (alternate) swaths 20-50 feet wide that are spaced at least one swath width apart.

Re-treat the same swath or bait previously untreated swaths if 75% or more of the amount originally placed has been consumed, if 0.4 inches of precipitation has fallen, or after 7 days of exposure to birds. Do not replenish bait if target species are no longer present or are no longer feeding on bait. Do not exceed the application rate limits specified below.

**Application Rates:** Regardless of the application method selected, do not make any treatment which exceeds rates of 58 lbs/treated acre of cracked corn or rolled barley baits, 110 lbs/treated acre of diluted poultry pellet bait, or 137 lbs/treated acre of diluted brown rice bait. Do not make more than 5 treatments per year to any one treated area.

## **DIRECTIONS FOR USE (Continued)**

### **POSTTREATMENT CLEANUP**

Beginning one day after bait application, search treated staging area, immediate surrounding areas, and known roost sites (if accessible). Collect carcasses of dead or dying birds and bury or burn them, according to applicable laws. Repeat carcass searches at 1-2 day intervals as long as bait is exposed and likely to remain toxic.

At the end of the baiting period, recover as much unconsumed bait material as possible and dispose of it in accordance with this label and applicable Federal, State and local laws.

### **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store only in original container, in a dry place inaccessible to children, pets, and domestic animals.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinseate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bags by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. If bags are not to be recycled, dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities. If burned, stay out of smoke.