

## RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

### SECTION 24(C) REGISTRATION FOR SPECIAL LOCAL NEED FOR SALE AND USE ONLY IN THE STATE OF KANSAS



All Applicable Directions, Restrictions and Precautions on the Registered Product Label for Stallion Insecticide (EPA Registration Number 279-9545) are to be followed.

These directions for use must be in the possession of the user at the time of application.

EPA SLN Number: KS - 110002

CROP	PEST	USE RATE
Alfalfa	Alfalfa Weevil Larva Alfalfa Weevil Adults Aphid species	9.25 – 11.75 fl. Oz/A

#### DIRECTIONS FOR USE:

##### Foliar Applications :

- Apply Stallion as a broadcast foliar spray with either ground or aerial application equipment. Use a minimum of 10 gpa by ground or 2 gpa by aerial application. Higher volumes of finished spray can improve insect control when temperatures are high, foliage is dense and/or when insect pressure is heavy.
- Stallion is labeled for Chemigation application. Refer to Chemigation Use Directions on the full Stallion label for details on this application.
- Make applications when insect populations reach economic thresholds or when significant feeding is occurring. Refer to local Cooperative Extension Pest Management Guidelines and/or scouting results.
- This product is highly toxic to bees exposed to direct treatment. Do not spray if bees are actively foraging in the area to be treated.

#### RESTRICTIONS:

- Observe all spray drift precautions on the EPA registered label.
- Applications may be made up to 7 days prior to cutting, grazing, or harvesting seed.
- Do not make applications of Stallion or follow with other products containing chlorpyrifos less than 10 days apart.
- Do not exceed 35.25 fl. Oz/A or make more than 4 applications of Stallion or other products containing chlorpyrifos per crop growing season.
- Do not exceed the maximum of 0.075 lb ai/acre of zeta-cypermethrin per crop growing season. This includes Stallion and other products containing zeta-cypermethrin.
- Re-entry Interval : 24 hours. Personal protective equipment is required for early entry.

#### Dealers Should Sell in Original Packages Only.

##### Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

